
XXIII. Descriptions of two new Species of Phalæna. By Mr. Louis Bosc, of Paris, Foreign Member of the Linnean Society.

Read October 5, 1790.

i. PYRALIS tuberculana.

P. ALIS anticis griseis fusco punctatis margine crassiori antice trituberculato.


H. Parisii. Larva in Hedyfaro Onobrichide et Coronillâ coronatâ.

Larva fusco viridis. *Folliculum* sportæforme, e parenchymâ plantarum fabricatum et cauli affixum. *Pupa* Junio occurrit, et *Imago* Aprili sequentis anni.

Caput ex albo argenteum; *Palpi* compressi, fuscæ, albo punctati; *Antennæ* fuscæ basi subtus argenteæ et auriculatæ; *Oculi* nigri. *Thorax* argenteo-griseoque varius, antice cristatus; *Crista* argentea, fusco-bifasciata; fascia anteriori minori. *Alæ* deflexæ; *Superiores* supra albæ, fusco griseoque punctatæ et maculatæ; *Tubercula* tria seriem formantia margine antice crassiori, omnia æqualia non scabra, dimidio alba et fusca. *Alæ* inferiores pallide fuscæ, puncto centrali nigro. *Pedes* fuscæ, albo annulati.

TAB. 17. Fig. 4. Pyralis tuberculana.

5. *Folliculus* pupam continens.



1. *Echeneis lineata.*




2. *Fasciola clavata.*



3. *Hirudo branchiata.*



Phalena Pyralis Tuberculana.



Phalena Tinea Sparrmannella.